



DNAbiotech

Biotechnology is our expertise

SDS-PAGE Gel preparation Kit

Catalog no.: DSK100

For preparation of about 40 gels

Related products:

Bradford Protein Assay Kit (cat#: DB0017)

Coomassie Brilliant Blue R-250 Staining Solution (cat#: DB9608)

Coomassie Brilliant Blue G-250 Staining Solution (cat#: DB9609)

Coomassie Brilliant Blue destaining Solution (cat#: DB9610)

DNAbiotech Co.

Intended for research use only

Modares Technology and Sciences Park, Room 507. No. 15, Gordafarid-Hayat junction, North Kargar,
Enghelan Square, Tehran, I.R. Iran

www.dnabiotch.ir

Cell phone: +989128382915

Introduction

The separation of macromolecules in an electric field is called *electrophoresis*. A very common method for separating proteins by electrophoresis uses a discontinuous polyacrylamide gel as a support medium and sodium dodecyl sulfate (SDS) to denature the proteins. The method is called sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE). SDS (also called lauryl sulfate) is an anionic detergent, meaning that when dissolved its molecules have a net negative charge within a wide pH range. A polypeptide chain binds amounts of SDS in proportion to its relative molecular mass. The negative charges on SDS destroy most of the complex structure of proteins, and are strongly attracted toward an anode (positively-charged electrode) in an electric field. Polyacrylamide gels restrain larger molecules from migrating as fast as smaller molecules. Because the charge-to-mass ratio is nearly the same among SDS-denatured polypeptides, the final separation of proteins is dependent almost entirely on the differences in relative molecular mass of polypeptides. In a gel of uniform density the relative migration distance of a protein (R_f , the f as a subscript) is negatively proportional to the log of its mass. If proteins of known mass are run simultaneously with the unknowns, the relationship between R_f and mass can be plotted, and the masses of unknown proteins estimated.

Protein separation by SDS-PAGE can be used to estimate relative molecular mass, to determine the relative abundance of major proteins in a sample, and to determine the distribution of proteins among fractions. The purity of protein samples can be assessed and the progress of a fractionation or purification procedure can be followed. Different staining methods can be used to detect rare proteins and to learn something about their biochemical properties. Specialized techniques such as Western blotting, two-dimensional electrophoresis, and peptide mapping can be used to detect extremely scarce gene products, to find similarities among them, and to detect and separate isoenzymes of proteins.

Storage Conditions and Reagent Preparation:

Kit will arrive packaged as a 4° C & will performed as specified if stored as directed and used before expiration date

Materials Supplied

No.	Item	QTY	Storage
1	Acrylamide/Bisacrylamide-stock solution (30:0~8 %)	100 mL	4° C
2	Separation gel buffer	80 mL	4° C
3	stacking gel buffer	25 mL	4° C
4	5 X sample buffer	2 × 6 mL	4° C
5	TEMED (100 %)	1 mL	4° C or RT
6*	Ammoniumpersulphate	10 Calibrated tubes	4° C or RT
7	1X Running buffer for 1 L	6 Packs (for 6 L)	4° C or RT

* Each tube contains 1 gr APS which should be reconstituted to 10% APS solution just before use. This Reagent must be used freshly. If any of the items listed above are missing or damaged, please contact us at www.dnabiotech.ir.

Reagent preparation

- Add 1 mL of DDW to each vial of APS. Each vial of **reconstituted APS** provides 1 mL of 10X (or 100%) APS Solution.
- Dissolve the ready to use powder of SDS-PAGE running buffer (cat#: DB0016) in 1 litre of DDW. This of reconstituted **1X** buffer can be used 2-3 times.

Procedure

WARNING: Unpolymerized acrylamide is a neurotoxin!

An intact SDS PAGE electrophoresis system should include: a tank, lid with power cables, electrode assembly, casting stands, casting frames, combs (usually 10-well or 15-well), and glass plates (thickness 0.75mm or 1.0mm or 1.5mm).

- (1) Clean the plates and combs. For each gel, you will need one short plate, one spacer plate, and one comb. These are usually found on the gray rack by the sink. Spray a little bit of 70% ethanol on the plates, and wipe dry with cotton. Wash the combs thoroughly with tap water. It is critical to remove all dust and small particles, especially any bits of left-over polyacrylamide.
- (2) Set-up the plates on the rack. Layer the short plate on the spacer plate, with the spacers in between and slide the two plates into the green holder. Make sure that the bottom edges of the two plates are flush to avoid leakage. Lock the plates in, and place the holder on the rack, with the bottom edges of the plates pushed into the gray foam gasket to make a water-tight seal. Test the seal by pipetting or squirting a small volume of water between the plates and making sure there is no leakage. Blot dry with filter paper.
- (3) **Pour the separating gel.** For each minigel (1 mm thick) you will need slightly more than 5 mL of reagent mix. Use the table below as a guide to calculate the total volumes you will need. Pipette solutions in order. Avoid introducing bubbles, which will inhibit polymerization. Swirl the

solution gently to mix thoroughly after addition of each component.

NOTE: The acrylamide percentage in SDS PAGE gel depends on the size of the target protein in the sample. (Details shown below)

Acrylamide %	M.W. Range
7%	50 kDa - 500 kDa
10%	20 kDa - 300 kDa
12%	10 kDa - 200 kDa
15%	3 kDa - 100 kDa

All given amounts are in mL

Final Acylamide percentage (%)	7.5	10	12.5	15	17.5
Separation gel buffer	1.5	1.5	1.5	1.5	1.5
Acrylamide/Bisacrylamide-stock solution	1.45	1.9	2.45	2.95	3.4
DDW	3.05	2.55	2.05	1.6	1.1
Ammoniumpersulphate (10 % or X) *	0.04	0.04	0.04	0.04	0.04
TEMED (15 %)*	0.05	0.05	0.05	0.05	0.05

- (4) Pour the stacking gel. For each gel you will need about 2 mL of reagent mix. Again, pipette the solutions carefully and swirl to mix after addition of each component. Pipette the gel mix between the plates up to just below the edge of the short plate.

All given amounts are in mL

Final Acylamide percentage (%)	3	4	5
Stacking gel buffer	0.625	0.625	0.625
Acrylamide/Bisacrylamide-stock solution	0.25	0.325	0.41
DDW	1.6	1.525	1.45
Ammoniumpersulphate (10 % or X) *	0.04	0.04	0.04
TEMED (15 %)*	0.05	0.05	0.05

* These reagents should be prepared freshly. You can either add from their stock solutions too, but the final amount should be optimized by the operator in the lab.


- **Once** TEMED is added, the gel will begin to polymerize, so you need to work fast (but carefully). Pipette the gel mix between the plates, making sure you leave enough space at the top for the stacking gel and comb. Carefully layer water on top of the gel solution.
- **Once** the gel has polymerized (about 10-15 mins), wash off the top of the gel with water. Carefully blot off excess water with a filter paper. Take care not to disturb or damage the top

of the gel.

- **Once** the gel has polymerized, slowly remove the comb under running water. Wash the wells carefully to avoid distorting them.

NOTE, Gel storage: Put back the comb, and sandwich the gel between two wet pieces of paper towels. The gel can be stored horizontally (with the cassette) at 4 °C for up to 1 week.

- (5) Carefully place in the comb like the following image.



- (6) According to your electrophoresis set assemble the cassette vertically.
- (7) Pour the running buffer (adjust the volume according to the size of your tank) and bring out the comb carefully.
- NOTE:** remove the bubbles formed at the bottom of the gel.
- (8) **Prepare the samples:** mix 1 volume of the sample and 4 volumes of the 5X sample loading buffer in appropriated microtube and incubate in boiling water for 5-10 minutes.
- (9) Spin the microtube containing prepared samples and load appropriated volume of them into the wells.
- (10) Apply electric current to the electrophoretic chamber by an authorized power supply. Set voltage to 100 - 150 volts (V) and current to 25 mA for one gel 80x80x1 mm, 12% PAA. Stop the run after blue front dye (Bromophenol-blue) has reached bottom of the gel (after approx. 1 - 1.5 hours).
- (11) Remove the gel from the electrophoretic chamber, disassemble the glass plate/ gel sandwich diligently, then fix and stain the gel for visualization of protein bands following appropriate protocols

Troubleshooting

Problem	Reason	Countermeasure
No current	Unclosed circuit	Check contacts/leads at source of current and separation chamber; check buffer level
Low current	Wrong adjustment of parameters at power source	For limiting amperage select the maximum voltage recommended for the chamber; for limiting voltage select maximum amperage
'Smile effect' at buffer front	Overheating	Pre-cool buffer; cooling via cooling circulator or a reduction in amperage
Slow migration of buffer front	Running buffer fully consumed	Always use fresh running buffer
Blurred bands	Diffusion after application of samples	Apply samples quickly; begin electrophoresis straight away
Blurred bands	Diffusion after separation	Transfer gel to fixing or staining solution immediately after electrophoresis
Irregular bands	Sample volumes too low or too different	Apply at least 5 µl sample; use approx. the same amounts of sample
Irregular bands	Differing saline content of samples	Desalinate samples as required (dialysis, gel filtration)
Formation of stripes	Precipitation of sample	Centrifuge or filter sample
Wide, partially smeared bands	Lipophilic substances in the sample	Remove substances prior to electrophoresis; increase SDS concentration if necessary
More bands than expected	Protease activity	Add protease inhibitor; minimize time between sample preparation and run
More bands than expected	Incomplete reduction	Check reduction conditions (if necessary prolong incubation time; increase DTT concentration)

DNAbiotech is the 1st Iranian Biotechnology Company producing wide ranges of ready to use **molecular buffers**. These products are produced from the high quality ingredients and help researchers run their projects faster.



Other services:

Gene, Primer and peptide synthesizing

- ✓ Cloning and expression of different recombinant peptides
- ✓ Bioinformatics services
- ✓ Production of different media and reagents in Cell and Bacteria culture
- ✓ Production of secondary antibodies (goat anti mouse and goat anti rabbit antibodies, HRP conjugated).
- ✓ And....

Our New Products Launch Coming Soon!

DNAbiotech Co.